

352-899

7/31/2014

1/22



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Registration
Number:

352-899

Date of Issuance:

JUL 3 1 2014

NOTICE OF PESTICIDE:
 Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance: **Conditional**

Name of Pesticide Product:
DuPont DPX-JMQ48 75XP Herbicide

Name and Address of Registrant (include ZIP Code):
E.I. du Pont de Nemours and Company
1007 Market Street
Wilmington, DE 19898

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA §3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration review of your product when the Agency requires all registrants of similar products to submit data. Data requirements can be found at www.regulations.gov. Thifensulfuron data requirements are outlined in GDCI-128845-1164 under docket ID EPA-HQ-OPP-2011-0171-0017. Tribenuron data requirements are outlined in GDCI-128887-974 under docket ID EPA-HQ-OPP-2010-0626-0017.
2. Submit Storage Stability (Guideline 830.6317) and Corrosion Characteristics (Guideline 830.6320) studies within eighteen (18) months from the date of this notice.

The basic confidential statement of formula (CSF) dated March 31, 2014 is acceptable.

A stamped copy of the label is enclosed for your records. Submit one (1) copy of the final printed label before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions regarding this Notice, please contact Mindy Ondish at (703)605-0723 or at ondish.mindy@epa.gov.

Signature of Approving Official:

Kable Bo Davis
Product Manager 25
Herbicide Branch
Registration Division (7505P)

Date:

JUL 3 1 2014



DuPont™ DPX-JMQ48 75XP

HERBICIDE


GROUP	2	HERBICIDE
-------	---	-----------

2/22

For Use on Wheat (including durum), Barley, Oat, Triticale and Fallow

Active Ingredients:	By Weight
Thifensulfuron-methyl	
Methyl 3-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl) amino]carbonyl]amino]sulfonyl]-2-thiophenecarboxylate	60%
Tribenuron methyl	
Methyl 2-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)methylamino]carbonyl]amino]sulfonyl]benzoate	15%
Other Ingredients:	25%
TOTAL	100%

EPA Reg. No. 352-899
Nonrefillable Container
 Net: _____
 OR
Refillable Container
 Net: _____



EPA Est. No. _____

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-441-3637 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution! Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.
 For medical emergencies involving this product, call toll free 1-800-441-3637.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves made out of any waterproof material.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Important: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "Applicators and Other Handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters or rinsate. Do not apply where/when conditions favor runoff.

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Assure accurate measurement of pesticides by all operation employees.
- Mix only enough product for the job at hand.
- Avoid over-filling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field/grove or mixing/loading station.
- Dilute and agitate excess solution and apply at labeled rates/uses.
- Avoid storage of pesticides near well sites.
- When triple rinsing the pesticide container, be sure to add the rinsate to the spray mix.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls.

Chemical-resistant gloves made out of any waterproof material.

Shoes plus socks.

DuPont™ DPX-JMQ48 75XP must be used only in accordance with instructions on this label, in separately issued labeling or exemptions under FIFRA (Supplemental Label, Special Local Need Registrations, FIFRA Section 18 exemptions, FIFRA 2(ee) bulletins), or as otherwise permitted by FIFRA. Always read the entire label, including the Limitation of Warranty and Liability.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specified by DuPont.

DPX-JMQ48 75XP is for use on wheat, barley, oat, triticale, post-harvest burndown, pre-plant burndown and fallow in most states. Check with your state extension service or Department of Agriculture before use, to be certain DPX-JMQ48 75XP is registered in your state.

PRODUCT INFORMATION

DPX-JMQ48 75XP herbicide is to be used in a tank mix with other suitable registered herbicides to provide selective postemergence control of certain broadleaf weeds in wheat (including durum), barley, oat, triticale, post-harvest burndown, preplant burndown and fallow. In the state of Arizona, DPX-JMQ48 75XP herbicide at 0.4 oz/a to 0.7 oz/a can be used alone or in a tankmix for control of broadleaf weeds in wheat, barley, oat and triticale.

DuPont™ DPX-JMQ48 75XP is a dispersible granule to be mixed in water or other recommended carrier and applied as a uniform broadcast spray. It is noncorrosive, nonflammable, nonvolatile and does not freeze.

RESTRICTIONS

Injury to or loss of adjacent sensitive crops, desirable trees or vegetation may result from failure to observe the following:

- Do not apply, drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants.
- Do not apply this product through any type of irrigation system.

Do not apply to wheat, barley, oat or triticale crops underseeded with another crop.

Do not harvest wheat, barley, oat or triticale sooner than 45 days after the last application of DPX-JMQ48 75XP.

When using DPX-JMQ48 75XP in tank mixes or sequential applications with other products containing thifensulfuron-methyl and/or tribenuron-methyl, do not exceed the following limits.

Use	Active Ingredient	Maximum oz ai per acre per Single Application	Maximum oz ai per acre per Use Period
wheat, barley triticale	thifensulfuron-methyl	0.45	0.75
	tribenuron-methyl	0.25	0.25
oats	thifensulfuron-methyl	0.3	0.3
	tribenuron-methyl	0.1	0.1
fallow, burndown, post harvest	thifensulfuron-methyl	0.45	0.75
	tribenuron-methyl	0.25	0.25

PRECAUTIONS

Injury to or loss of adjacent sensitive crops, desirable trees or vegetation may result from failure to observe the following:

- Take all necessary precautions to avoid all direct or indirect contact (such as spray drift) with non-target plants or areas.

Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than wheat, barley, oat or triticale.

Dry, dusty field conditions may result in reduced control in wheel track areas.

DPX-JMQ48 75XP should not be applied to wheat, barley, oat or triticale that is stressed by severe weather conditions, drought (including low levels of subsoil moisture), low fertility, water-saturated soil, disease, or insect damage, as crop injury may result. Risk of injury is greatest when crop is in the 2 to 5-leaf stage. Severe winter stress, drought, disease, or insect damage following application also may result in crop injury.

Wheat, barley, oat and triticale may differ in their response to various herbicides. DuPont recommends that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of DPX-JMQ48 75XP to a small area.

Under certain conditions, such as heavy rainfall, prolonged cold weather (daily high temperature less than 50°F), or wide fluctuations in day/night temperatures prior to or soon after DPX-JMQ48 75XP application, temporary discoloration and/or crop injury may occur. To reduce the potential of crop injury, tank mix DPX-JMQ48 75XP with 2,4-D (ester formulations perform best—see "Tank Mixtures" section of this label) and apply after the crop is in the tillering stage of growth.

BIOLOGICAL ACTIVITY AND ENVIRONMENTAL CONDITIONS

Best results are obtained when DPX-JMQ48 75XP is applied to young, actively growing weeds. The use rate will depend on weed spectrum and size of weed at time of application. The degree of control and duration of effect are dependent on rate used, sensitivity and size of target weed and environmental conditions at the time of and following application. DPX-JMQ48 75XP stops growth of susceptible weeds rapidly. However, typical symptoms of dying weeds (discoloration) may not be noticeable for 1-3 weeks after application (2-5 weeks for wild garlic, when present) depending on the environmental conditions and weed susceptibility. Warm, moist conditions following treatment promote the activity of DPX-JMQ48 75XP, while cold, dry conditions delay the activity. Weeds hardened-off by cold weather or drought stress will be less susceptible.

A vigorous growing crop will aid weed control by shading and providing competition for weeds. However, a dense crop canopy at time of application can intercept spray and result in reduced weed control. Weeds may not be adequately controlled in areas of thin crop stand or seeding skips.

Applications made to weeds that are in the cotyledon stage, larger than the size indicated, or to weeds under stress may result in unsatisfactory control.

DPX-JMQ48 75XP may injure crops that are stressed from adverse environmental conditions (such as extreme temperatures or moisture), abnormal soil conditions, or cultural practices. In addition, different varieties of the crop may have differing levels of sensitivity to treatment with DPX-JMQ48 75XP under otherwise normal conditions.

5/22

Treatment of sensitive crop varieties may injure crops. To reduce the potential of crop injury, tank mix DuPont™ DPX-JMQ48 75XP with 2,4-D (ester formulations perform best – see "TANK MIXTURES" section of this label) and apply after the crop is in the tillering stage of growth.

Weed control may be reduced if rainfall or snowfall occurs soon after application. Several hours of dry weather are needed to allow DPX-JMQ48 75XP to be sufficiently absorbed by weed foliage.

RESISTANCE

DPX-JMQ48 75XP, which contains the active ingredients thifensulfuron methyl and tribenuron methyl, is a Group 2 herbicide based on the mode of action classification system of the Weed Science Society of America.

When herbicides with mode of action classifications that affect the same biological sites of action are used repeatedly over several years to control the same weed species in the same treatment area, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that area. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different biological site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes; it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that affect a different site of action. Weed escapes that are allowed to go to seed and movement of plant material between treatment areas on equipment will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative to determine appropriate actions for treating specific resistant weed biotypes in your area.

INTEGRATED PEST MANAGEMENT

DuPont recommends the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an IPM program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

LABELLED USES

DPX-JMQ48 75XP herbicide provides selective postemergence control of certain broadleaf weeds in wheat (including durum), barley, oat, triticale, post-harvest burndown, pre-plant burndown and fallow.

APPLICATION TIMING

Wheat (Including Durum), Barley, Winter Oat and Triticale

Make applications after the crop is in the 2-leaf stage, but before the flag leaf is visible. Do not harvest within 45 days of the last application.

Spring Oat

Make applications after the crop is in the 3-leaf stage but before jointing. Do not use on "Ogle", "Porter" or "Premier" varieties as crop injury can occur. Do not harvest within 45 days of the last application.

Pre-Plant Burndown

For burndown of emerged weeds, broadcast applications of DPX-JMQ48 75XP may be applied up through planting, but before wheat (including durum), barley, or triticale plants emerge. DPX-JMQ48 75XP can be used as a burndown treatment prior to planting other crops. See "CROP ROTATION" for the time interval required before planting.

Post Harvest

DPX-JMQ48 75XP may be used as a burndown treatment to crop stubble when the majority of weeds have emerged and are actively growing. (See the "CROP ROTATION" section of this label for additional information).

Fallow

Apply DPX-JMQ48 75XP in the spring or fall when the majority of weeds have emerged and are actively growing. Generally, such applications are made in the spring or fall when most cereal applications are made. (See the "CROP ROTATION" section of this label for additional information).

USE RATES

Unless otherwise specified by DuPont, do not use less than 0.4 ounce DuPont™ DPX-JMQ48 75XP per acre.

Wheat (including Durum), Barley and Triticale

Apply 0.4 to 0.7 ounce DPX-JMQ48 75XP per acre.

Sequential treatments of DPX-JMQ48 75XP may be made provided the total amount of DPX-JMQ48 75XP applied to the crop does not exceed 1.2 ounces per acre.

Oat (Spring and Winter)

Apply 0.4 to 0.5 ounce DPX-JMQ48 75XP per acre. Do not make more than one application of DPX-JMQ48 75XP per crop season on oat.

Pre-Plant Burndown

Apply 0.4 to 0.7 ounce DPX-JMQ48 75XP per acre as a burndown treatment prior to planting any crop; or shortly after planting, but prior to emergence of, wheat (including durum), barley, or triticale. See "CROP ROTATION" for the time interval required before planting.

DPX-JMQ48 75XP should be applied in combination with other suitable registered preplant burndown herbicides (See the "TANK MIXTURES" section of this label for additional information).

Sequential treatments of DPX-JMQ48 75XP may also be made provided the total amount of DPX-JMQ48 75XP applied during one fallow/preplant season does not exceed 1.2 ounces per acre.

Post Harvest and Fallow

Apply 0.4 to 0.7 ounce DPX-JMQ48 75XP per acre as a postemergence fallow treatment, in combination with other suitable registered fallow herbicides (See the "TANK MIXTURES" section of this label for additional information). See "CROP ROTATION" for the time interval required before planting.

Sequential treatments of DPX-JMQ48 75XP may be made provided the total amount of DPX-JMQ48 75XP applied in fallow does not exceed 1.2 ounces per acre.

SPRAY ADJUVANTS

Include a spray adjuvant with applications of DPX-JMQ48 75XP. An ammonium nitrogen fertilizer may also be used. Do not use low rates of liquid nitrogen fertilizer solution as a substitute for a surfactant. Always use a surfactant, unless otherwise recommended. Antifoaming agents may be used if needed.

Consult your Ag dealer or applicator, local DuPont fact sheets and technical bulletins prior to using an adjuvant system. Select adjuvants that are authorized for use with all products in an DPX-JMQ48 75XP tank mix. Products must contain only EPA-exempt ingredients.

Nonionic Surfactant (NIS)

- Apply 0.25 to 0.50% volume/volume (2 pints to 4 pints per 100 gal of spray solution).
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12. – See the "TANK MIXTURES" section of this label for additional information.

Crop Oil Concentrate (COC) - Petroleum or Modified Seed Oil (MSO)

- Apply at least 1% v/v (1 gal per 100 gal spray solution), or 2% under arid conditions. MSO adjuvants may be used at 0.5% v/v if specified on local DuPont product literature or service policies.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

Special Adjuvant Types

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality and have been evaluated and approved by DuPont product management. Consult separate DuPont technical bulletins for detailed information before using adjuvant types not specified on this label.

Ammonium Nitrogen Fertilizer

- Use 2 qt/acre of a high-quality urea ammonium nitrate (UAN) with a surfactant, such as 28%N or 32%N, or 2 lb/acre of a spray-grade ammonium sulfate (AMS), with a surfactant. Use 4 qt/acre UAN or 4 lb/acre AMS under arid conditions.
- See TANK MIXTURES With Liquid Nitrogen Solution Fertilizers for instructions on using fertilizer as a carrier in place of water.

7/22

WEEDS CONTROLLED WHEN TANK-MIXED WITH BROMOXYNIL CONTAINING PRODUCTS

(Such as "Buctril", "Bison", "Bronate" or "Bronate Advanced" or "Rhino")

- | | |
|-------------------------------------|--------------------------------|
| Annual knawel | Marshelder |
| Annual sowthistle | Miners lettuce |
| Black mustard | Mouseear chickweed |
| Black nightshade | Pennsylvania smartweed |
| Bushy wallflower/Treacle
mustard | Pepperweed species |
| Carolina geranium | Prickly lettuce*‡ |
| Coast fiddleneck | Prostrate knotweed |
| Common buckwheat | Puncturevine |
| Common chickweed* | Redmaids |
| Common cocklebur | Redroot pigweed |
| Common groundsel | Redstem filaree |
| Common lambsquarters | Russian thistle*‡ |
| Common ragweed | Scentless
chamomile/mayweed |
| Common sunflower* | Shepherd's-purse |
| Common tarweed | Silverleaf nightshade |
| Corn chamomile | Smallflower buttercup |
| Corn gromwell | Smooth Pigweed |
| Corn spurry | Spiny pigweed |
| Cow cockle | Stinking
mayweed/Dogfennel |
| Cress (mouse-ear) | Swinecress |
| Cutleaf nightshade | Tall morningglory |
| Curly dock | Tall waterhemp |
| Eastern black nightshade | Tansymustard |
| False chamomile | Tartary buckwheat |
| Field pennycress | Tarweed fiddleneck |
| Flixweed | Tumble/Jim Hill mustard |
| Fumitory | Velvetleaf |
| Giant Ragweed | Volunteer canola |
| Green smartweed | Volunteer lentils |
| Hemp sesbania | Volunteer peas |
| Henbit | Volunteer sunflower* |
| Horned poppy | White cockle |
| Ivyleaf morningglory | Wild buckwheat |
| Jimsonweed | Wild chamomile |
| Kochia *‡ | Wild mustard |
| Ladysthumb | Wild radish |
| Lanceleaf sage | Yellow rocket |
| London rocket | |
| Mallow (little) | |

PARTIAL CONTROL**

- | | |
|----------------|-------------------------|
| Canada thistle | Cutleaf eveningprimrose |
| Common mallow | Marestail |

* See SPECIFIC WEED INSTRUCTIONS for more information.

**Partial control: A visual reduction of weed population as well as a significant loss of vigor for individual weed plants. For better results, use 6 ounce active ingredient per acre of bromoxynil containing herbicide (such as "Bronate" or "Bison" at 1 1/2 pint per acre - refer to the "USE RATES" section of this label).

‡ Naturally occurring resistant biotypes of kochia, prickly lettuce, and Russian thistle are known to occur. See the "TANK MIXTURES" and "SPECIFIC WEED INSTRUCTIONS" sections of this label for additional details.

WEEDS CONTROLLED WHEN TANK-MIXED WITH 2,4-D CONTAINING PRODUCTS

(Such as "Agri-Star", "Barrage", "Omni-Amine" or "Weedar 64")

Annual knawel	Miners lettuce
Annual sowthistle	Mouseear chickweed
Black mustard	Pennsylvania smartweed
Bushy wallflower/Treacle mustard	Pepperweed species
Carolina geranium	Prickly lettuce*‡
Coast fiddleneck	Prostrate knotweed
Common buckwheat	Puncturevine
Common cocklebur	Redmaids
Common groundsel	Redroot pigweed
Common lambsquarters	Redstem filaree
Common mallow	Russian thistle*‡
Common purselane	Scentless chamomile/mayweed
Common sunflower*	Shepherd's-purse
Common ragweed	Smallflower buttercup
Common tarweed	Smooth Pigweed
Corn chamomile	Spiny pigweed
Corn spurry	Stinking mayweed/Dogfennel
Cow cockle	Swinecress
Cress (mouse-ear)	Tansymustard
Cutleaf nightshade	Tarweed fiddleneck
Curly dock	Tumble/Jim Hill mustard
False chamomile	Velvetleaf
Field pennycress	Volunteer canola
Flixweed	Volunteer lentils
Giant ragweed	Volunteer peas
Green smartweed	Volunteer sunflower*
Henbit	White cockle
Ivyleaf-morningglory	Wild buckwheat
Kochia *‡	Wild chamomile
Ladysthumb	Wild mustard
London rocket	Wild radish
Mallow (little)	
Marshelder	

PARTIAL CONTROL**

Canada thistle	Marestail
Corn gromwell	Tall morningglory
Fumitory	Tall waterhemp
Hemp sesbania	

* See SPECIFIC WEED INSTRUCTIONS for more information.

**Partial control: A visual reduction of weed population as well as a significant loss of vigor for individual weed plants. For better results, use higher rates 2,4-D containing herbicides (such as "Barrage" or "AgriStar" - refer to the "USE RATES" sections of these labels).

‡ Naturally occurring resistant biotypes of kochia, prickly lettuce, and Russian thistle are known to occur. See the "TANK MIXTURES" and "SPECIFIC WEED INSTRUCTIONS" sections of this label for additional details.

9/22

WEEDS CONTROLLED WHEN TANK-MIXED WITH 2,4-D + DICAMBA CONTAINING PRODUCTS

(Such as "Banvel", "Banvel + 2,4-D or "Clarity")

Annual knawel
 Annual sowthistle
 Black mustard
 Bushy wallflower/Treacle
 mustard
 Carolina geranium
 Coast fiddleneck
 Common buckwheat
 Common cocklebur
 Common groundsel
 Common lambsquarters
 Common mallow
 Common purselane
 Common sunflower*
 Common ragweed
 Common tarweed
 Corn chamomile
 Corn spurry
 Cow cockle
 Cress (mouse-ear)
 Cutleaf nightshade
 Curly dock
 False chamomile
 Field pennycress
 Flixweed
 Fumitory
 Giant ragweed
 Green smartweed
 Hemp sesbania
 Henbit
 Ivyleaf morningglory
 Kochia *‡
 Ladysthumb
 London rocket
 Mallow (little)
 Marshelder

Miners lettuce
 Mouseear chickweed
 Pennsylvania smartweed
 Pepperweed species
 Prickly lettuce*‡
 Prostrate knotweed
 Puncturevine
 Redmaids
 Redroot pigweed
 Redstem filaree
 Russian thistle*‡
 Scentless
 chamomile/mayweed
 Shepherd's-purse
 Smallflower buttercup
 Smooth Pigweed
 Spiny Pigweed
 Stinking
 mayweed/Dogfennel
 Swinecress
 Tall morningglory
 Tall waterhemp
 Tansymustard
 Tarweed fiddleneck
 Tumble/Jim Hill mustard
 Velvetleaf
 Volunteer canola
 Volunteer lentils
 Volunteer peas
 Volunteer sunflower*
 White cockle
 Wild buckwheat
 Wild chamomile
 Wild mustard
 Wild radish

PARTIAL CONTROL**

Canada thistle
 Corn gromwell

Marestail
 Spiny pigweed

* See SPECIFIC WEED INSTRUCTIONS for more information.

**Partial control: A visual reduction of weed population as well as a significant loss of vigor for individual weed plants. For better results, use higher rates 2,4-D and/or dicamba containing herbicides (such as "Barrage", "AgriStar", "Banvel", "Banvel SFG" or "Clarity" - refer to the "USE RATES" sections of these labels).

‡ Naturally occurring resistant biotypes of kochia, prickly lettuce, and Russian thistle are known to occur. See the "TANK MIXTURES" and "SPECIFIC WEED INSTRUCTIONS" sections of this label for additional details.

10/22

WEEDS CONTROLLED WHEN TANK-MIXED WITH FLUROXYPYR CONTAINING PRODUCTS

(Such as "Starane", "Starane" Ultra, "Starane" NXT, "Starane +Saber", "Starane +Sword" or "Starane +Salvo")

- | | |
|-------------------------------------|--------------------------------|
| Annual knawel | Morningglory species *** |
| Annual sowthistle | Mouseear chickweed |
| Bedstraw (cleavers) *** | Pennsylvania smartweed |
| Black mustard | Prickly lettuce *** ‡ |
| Bushy wallflower/Treacle
mustard | Prostrate knotweed |
| Carolina geranium | Puncturevine *** |
| Coast fiddleneck | Redmaids |
| Coffeeweed *** | Redroot pigweed |
| Common buckwheat | Redstem filaree |
| Common chickweed *** | Russian thistle * ‡ |
| Common cocklebur *** | Scentless
chamomile/mayweed |
| Common groundsel | Shepherd's-purse |
| Common lambsquarters | Smallflower buttercup |
| Common purslane *** | Stinking
mayweed/Dogfennel |
| Common ragweed *** | Swinecress |
| Common sunflower *** | Tansymustard |
| Corn chamomile | Tarweed fiddleneck |
| Corn spurry | Tumble/Jim Hill mustard |
| Cress (mouse-ear) | Velvetleaf *** |
| Curly dock | Venice mallow *** |
| False chamomile | Volunteer canola |
| Field pennycress | Volunteer flax *** |
| Flixweed | Volunteer lentils |
| Green smartweed | Volunteer peas |
| Hemp dogbane *** | Volunteer sunflower * |
| Kochia * ‡ | White cockle |
| Ladysthumb | Wild-buckwheat |
| London-rocket | Wild chamomile |
| Mallow (little) | Wild mustard |
| Marshelder | White clover *** |
| Miners lettuce | |

PARTIAL CONTROL**

- | | |
|--------------------------|-----------------------|
| Black nightshade | Field horsetail |
| Canada thistle | Henbit |
| Common mallow | Marestail |
| Cutleaf nightshade | Silverleaf nightshade |
| Eastern black nightshade | Volunteer potato § |
| Field Bindweed | |

* See SPECIFIC WEED INSTRUCTIONS for more information.

**Partial control: A visual reduction of weed population as well as a significant loss of vigor for individual weed plants. Use 1 1/2 - 2 ounce active ingredient per acre of fluroxypyr containing herbicide (such as "Starane" at 1/2 - 2/3 pint per acre - refer to the "USE RATES" section of this label).

*** Use 1 1/2 - 2 ounce active ingredient per acre fluroxypyr containing herbicides (such as "Starane" at 1/2 - 2/3 pint per acre).

‡ Naturally occurring resistant biotypes of kochia, prickly lettuce and Russian thistle are known to occur. See the "TANK MIXTURES" and "SPECIFIC WEED INSTRUCTIONS" sections of this label for additional details.

§ Use 2-4 ounce active ingredient per acre fluroxypyr containing herbicides (such as "Starane" at 1 1/3 pint per acre). See specific fluroxypyr containing herbicide label for rate recommendation and precautions.

SPECIFIC WEED INSTRUCTIONS

Common chickweed: For best results, apply DuPont™ DPX-JMQ48 75XP in a tank mix with either bromoxynil or fluroxypyr when all or the majority of weeds have germinated and are past the cotyledon stage and less than 3 inches tall or across. When mixing with bromoxynil, use a minimum of 6 ounces active ingredient per acre (such as "Bronate" or "Bison" at 1 1/2 pint per acre). When mixing with fluroxypyr, use a minimum of 1 1/2 ounces active ingredient per acre (such as "Starane" at 1/2 pint per acre).

Kochia: Naturally occurring biotypes resistant to DPX-JMQ48 75XP are known to occur.

For best results, DPX-JMQ48 75XP in a tank mix with CleanWave, WideMatch, Colt, or herbicides containing the active ingredient bromoxynil or fluroxypyr. See "TANK MIXTURES" for additional information.

Prickly lettuce: Naturally occurring biotypes resistant to DPX-JMQ48 75XP are known to occur. For best results, DPX-JMQ48 75XP tank mixed with a minimum of 1 1/2 ounces active ingredient per acre of fluroxypyr containing herbicide (such as "Starane" at 1/2 pint per acre) should be applied in the spring when prickly lettuce are 2" to 4" across and are actively growing.

Russian Thistle: Naturally occurring biotypes resistant to DPX-JMQ48 75XP are known to occur. DPX-JMQ48 75XP should be applied in the spring when Russian thistle are less than 2" tall and are actively growing. Apply a minimum of 6 ounces active ingredient per acre of a bromoxynil containing herbicide (such as "Bronate" or "Bison" at 1 1/2 pints per acre) when all or the majority of weeds have germinated.

DPX-JMQ48 75XP can also be tank mixed with a minimum of 1 1/2 ounces active ingredient per acre of a fluroxypyr and 2,4-D or MCP containing herbicide (such as "Starane +Saber" at 1 1/2 pints per acre, "Starane +Sword" at 1 1/8 pints per acre or "Starane +Salvo" at 1 pint per acre) and should be applied in the spring when Russian thistle are less than 2" tall and are actively growing.

SU / Clearfield Tolerant Volunteer Sunflowers: For suppression, apply a minimum of 1 1/2 ounces active ingredient per acre of a fluroxypyr containing herbicide (such as "Starane" at 1/2 pint per acre).

For improved results, apply a minimum of 6 ounces active ingredient per acre of a bromoxynil containing herbicide (such as "Bronate" or "Bison" at 1 1/2 pints per acre). Delay application until first sunflower seedlings emerging are 4 inches in height.

For improved results, DPX-JMQ48 75XP tank mixed with a minimum of 1 1/2 ounces active ingredient per acre of a fluroxypyr and 2,4-D or MCP containing herbicide (such as "Starane +Saber" at 1 1/2 pints per acre, "Starane +Sword" at 1 1/8 pints per acre or "Starane +Salvo" at 1 pint per acre) should be applied in the spring when SU/Clearfield tolerant volunteer sunflower are less than 2" tall and are actively growing.

TANK MIXTURES IN CEREALS

Read and follow all manufacturers' label instructions for any companion herbicides, fungicides, and/or insecticides. If those instructions conflict with this label, do not tank mix that product with DPX-JMQ48 75XP. Read and follow all label instructions on timing, precautions, and warnings for any companion products before using these tank mixtures. Follow the most restrictive labeling.

2,4-D (amine or ester) or MCPA (amine or ester)

DPX-JMQ48 75XP may be tank mixed with the amine and ester formulations of 2,4-D or MCPA herbicides for use on wheat, barley, or fallow (MCP or MCPA can also be used for oat).

For best results in the Red River Valley and adjacent areas of North Dakota and Minnesota, add the ester formulations of 2,4-D or MCPA herbicides to the tank at 3/8 lb active ingredient (such as 3/4 pint of a 4 lb/gal product, 1/2 pint of a 6 lb/gal product). No additional surfactant is needed with this mixture.

For best results, in other areas, add the ester formulations of 2,4-D or MCPA herbicides to the tank at 1/4 to 3/8 lb active ingredient (such as 1/2 to 3/4 pint of a 4 lb/gal product, 1/3 to 1/2 pint of a 6 lb/gal product). Nonionic surfactant may be added to the mixture at 1/2 to 1 quart per 100 gal of spray solution (0.125 to 0.25% v/v); however, adding nonionic surfactant may increase the potential for crop injury, especially at the higher phenoxy rates. Higher rates of 2,4-D or MCPA may be used, but do not exceed the highest rate allowed by those respective labels.

With dicamba (such as "Banvel"/"Banvel" SGF/"Clarity")

DPX-JMQ48 75XP may be tank mixed with 1/16 to 1/8 lb active ingredient dicamba (such as 2-4 fluid ounces of "Banvel", 4-8 fluid ounces of "Banvel" SGF, or 2-4 fluid ounces of "Clarity"). Use higher rates when weed infestation is heavy. Nonionic surfactant may be added to the mixture at 1/2 to 1 quart per 100 gal of spray solution (0.125 to 0.25% v/v); however, adding nonionic surfactant may increase the potential for crop injury. Refer to the specific dicamba label for application timing and restrictions. Tank mixes of DPX-JMQ48 75XP plus dicamba may result in reduced control of some broadleaf weeds.

With 2,4-D or MCP (amine or ester) and "Banvel"/"Clarity"

DPX-JMQ48 75XP may be applied in a 3-way tank mix with formulations of dicamba and 2,4-D or MCP. Make application of DPX-JMQ48 75XP plus 1/16 to 1/8 lb active ingredient dicamba (such as 2 to 4 fluid ounces of "Banvel", 4 to 8 fluid ounces of "Banvel" SGF, or 2 to 4 fluid ounces of "Clarity") plus 1/4 to 3/8 lb active ingredient 2,4-D or MCP ester or amine per acre. Use higher rates when weed infestation is heavy. Nonionic surfactant may be added to the mixture at 1/2 to 1 quart per 100 gal of spray

solution (0.125 to 0.25% v/v); however, adding nonionic surfactant may increase the potential for crop injury. Apply this three-way combination to winter wheat after the crop is tillering and prior to jointing (first node).

In Spring Wheat (including Durum), apply after the crop is tillering and before it exceeds the 5-leaf stage.

In Spring Barley, apply after the crop is tillering and before it exceeds the 4-leaf stage.

With Bromoxynil containing products (such as "Buctril", "Bison", "Bronate", "Bronate Advanced" or "Rhino")

DuPont™ DPX-JMQ48 75XP may be tank mixed with bromoxynil containing herbicides registered for use on wheat, barley or triticale. For best results, add bromoxynil containing herbicides to the tank at 6 to 12 oz active ingredient per acre (such as "Bronate" or "Bison" at 3/4 to 1 1/2 pt per acre). Tank mixes of DPX-JMQ48 75XP plus bromoxynil may result in reduced control of Canada thistle.

With fluroxypyr containing products (such as "Starane" brands)

DPX-JMQ48 75XP may be tank mixed with fluroxypyr containing herbicides for improved control of Kochia (2-4" tall) and other broadleaf weeds. For best results, add fluroxypyr containing herbicides to the tank at 1 to 2 oz active ingredient per acre (such as "Starane" 1/3 to 2/3 pints per acre). 2,4-D and MCP herbicides (preferably ester formulations) may be tank mixed with DPX-JMQ48 75XP plus fluroxypyr.

DPX-JMQ48 75XP may be used in combination with "Starane" NXT at 10 to 14 fluid ounces per acre for improved control of kochia less than 2" tall or at 14 to 21 fluid ounces per acre for kochia 2 to 4" tall.

With "Huskie" or "Wolverine" herbicides

DPX-JMQ48 75XP at 0.4 to 0.7 oz/a can be tank mixed with "Huskie" at 8.5 fl oz/a or "Wolverine" at 20 fl oz/a in wheat, durum, or barley for control of broadleaf weeds, including kochia (less than 2" in height). For larger weeds, higher labelled rates of "Huskie" or "Wolverine" are recommended.

With "CleanWave" Herbicide

For improved control of kochia and other broadleaf weeds in wheat (including durum), DPX-JMQ48 75XP may be tank mixed with "CleanWave". Tank mix "CleanWave" at 7 to 14 fluid ounces per acre for kochia less than 2" tall and at 14 ounces per acre for kochia 2 - 8" tall. Add 1 to 2 pints NIS per 100 gallons of spray solution in tank mixes of "CleanWave" with DPX-JMQ48 75XP (see SPRAY ADJUVANTS).

With "WideMatch" or "Colt" herbicides

For improved control of kochia, Canada thistle and other broadleaf weeds in wheat (including durum), barley, and oat, DPX-JMQ48 75XP may be tank mixed with "WideMatch" or "Colt". Tank mix at 1/2 to 2/3 pints per acre for kochia less than 2" tall and 2/3 to 1 pint per acre for kochia 2 - 4" tall. Add 1 to 2 pints NIS per 100 gallons of spray solution in tank mixes of WideMatch or Colt with DPX-JMQ48 75XP (see SPRAY ADJUVANTS).

With "Maverick"

DPX-JMQ48 75XP can be tank mixed with "Maverick" herbicide for improved control of grassy weeds in wheat.

DPX-JMQ48 75XP and a bromoxynil containing herbicide (such as "Bronate" or "Bison" at 3/4 to 1 pint per acre) may be tank mixed with 2/3 ounce per acre of "Maverick" herbicide for control of grassy weeds in wheat. This tank mix may also include "Starane" for greater spectrum of broadleaf control - see the "Maverick" label for specific use directions and restrictions. Apply 0.5% volume/volume (4 pint per 100 gal of spray solution) of non-ionic surfactant (NIS) with this tank mix. Some reduction in annual grass control may occur when optimum environmental conditions do not occur for several days prior to and after application - such as low moisture conditions, high and low temperatures, low humidity.

DPX-JMQ48 75XP and a fluroxypyr containing herbicide (such as "Starane", "Starane +Saber", "Starane +Sword" or "Starane +Salvo") may be tankmixed with 2/3 ounce per acre of "Maverick" herbicide for control of grassy weeds in wheat. Tank mixtures with herbicides formulated as amines may decrease the effectiveness of "Maverick" herbicide. Apply 0.5% volume/volume (4 pint per 100 gal of spray solution) of non-ionic surfactant (NIS) with this tankmix. Some reduction in annual grass control may occur when optimum environmental conditions do not occur for several days prior to and after application - such as low moisture conditions, high and low temperatures, low humidity.

With "Aim"

DPX-JMQ48 75XP can be tank mixed with "Aim" herbicide for improved control of weeds in wheat and barley.

With "Stinger", "Curtail" or "Curtail M"

DPX-JMQ48 75XP can be tank mixed with "Stinger", "Curtail" or "Curtail M" herbicide for improved control of weeds in wheat and barley.

DPX-JMQ48 75XP and fluroxypyr containing herbicides (such as "Starane", "Starane +Saber", "Starane +Sword" or "Starane +Salvo") may be tankmixed with "Stinger" or "Curtail" herbicide for improved control of weeds in wheat and barley.

With "Assert" Herbicide

DPX-JMQ48 75XP can be tank mixed with "Assert". When tank mixing DPX-JMQ48 75XP with "Assert", always include another broadleaf weed herbicide with a different mode of action (for example 2,4-D ester, MCP ester, or bromoxynil - such as "Buctril",

"Bison", "Bronate" or "Bronate Advanced"). Applications of DuPont™ DPX-JMQ48 75XP plus "Assert" may cause temporary crop discoloration, stunting, or injury when heavy rainfall occurs shortly after application.

DPX-JMQ48 75XP and fluroxypyr containing herbicides (such as "Starane", "Starane +Sword" or "Starane +Salvo") may be tank mixed with "Assert". Applications of DPX-JMQ48 75XP plus "Assert" may cause temporary crop discoloration, stunting, or injury when heavy rainfall occurs shortly after application.

Refer to the "Assert" label for specific instructions and restrictions when using amine formulations or additional tank mix products.

With "Axial"

For improved control of wild oats and other grasses, DPX-JMQ48 75XP at 0.4 to 0.7 ounces per acre may be tank mixed with "Axial" branded products in wheat and barley. Refer to Axial label for specific adjuvant recommendations.

With "Discover"

DPX-JMQ48 75XP can be tank mixed with "Discover" herbicide for improved control of grass weeds in spring wheat.

DPX-JMQ48 75XP and a bromoxynil containing herbicide (such as "Bronate" or "Bison" at 3/4 to 1 pint per acre) may be tank mixed with 4.0 ounces per acre of "Discover" herbicide, or 16 fluid ounces per acre "Discover" NG, for control of wild oat in wheat. This tank mix may also include "Starane" for greater spectrum of broadleaf control - see the "Discover" label for specific use directions, tank mixes, precautions, restrictions and geographical limitations of use.

DPX-JMQ48 75XP and a fluroxypyr containing herbicide (such as "Starane" or "Starane +Sword") may be tank mixed with 4.0 ounces per acre of "Discover" herbicide, or 16 fluid ounces per acre of "Discover" NG, for control of wild oat in wheat. See the "Discover" label for specific use directions, tank mixes, precautions, restrictions and geographical limitations of use. Some reduction in annual grass control may occur when optimum environmental conditions do not occur for several days prior to and after application - such as low moisture conditions, high and low temperatures or low humidity.

With "Everest"

DPX-JMQ48 75XP can be tank mixed with "Everest" herbicide for improved control of grassy weeds in spring wheat. When DPX-JMQ48 75XP and Everest are tank mixed, the mix must include 1/4 pint 2,4-D.

DPX-JMQ48 75XP and a bromoxynil containing herbicide (such as "Bronate" or "Bison" at 3/4 to 1 pint per acre) may be tank mixed with 0.3 ounce per acre of "Everest" for control of green foxtail, or 0.61 ounce per acre of "Everest" for control of green foxtail, yellow foxtail and wild oat. This tank mix may also include "Starane" for greater spectrum of broadleaf control - see the "Everest" label for specific use directions and restrictions.

DPX-JMQ48 75XP and a fluroxypyr containing herbicide (such as "Starane", "Starane +Saber", "Starane +Sword" or "Starane +Salvo") may be tankmixed with 0.3 ounce per acre of "Everest" for control of green foxtail or 0.61 ounce per acre of "Everest" for control of green foxtail, yellow foxtail and wild oat. See the "Everest" label for specific use directions, tankmixes, precautions and restrictions of use. Some reduction in annual grass control may occur when optimum environmental conditions do not occur for several days prior to and after application - such as low moisture conditions, high and low temperatures or low humidity.

With "Hoelon"

A tank mix of "Hoelon" 3EC herbicide + DPX-JMQ48 75XP herbicide can be applied for annual ryegrass (in the Pacific Northwest only), wild oat and broadleaf weed control in winter and spring wheat, and spring barley. The "Hoelon" 3EC herbicide rate should be 2 2/3 pints per acre with 0.4 ounce per acre of DPX-JMQ48 75XP herbicide in spring and winter wheat.

A three-way tank mix of "Hoelon" 3EC herbicide + "Buctril" herbicide + DPX-JMQ48 75XP herbicide can be applied for annual ryegrass (in the Pacific Northwest only), wild oat and broadleaf weed control in winter and spring wheat, and spring barley. The "Hoelon" 3EC herbicide rate should be 2 2/3 pints per acre with 0.4 ounce per acre DPX-JMQ48 75XP herbicide in winter wheat, spring wheat and spring barley. "Buctril" herbicide should be used at 1 pint per acre.

This tank mixture should only be used under good soil moisture conditions when wild oats are in the 1 to 4 leaf stage. Reduced control of foxtail is likely when tank mixing "Hoelon" with DPX-JMQ48 75XP herbicide. When foxtail is the major grassy weed in the field, DO NOT tank mix "Hoelon" 3EC herbicide + DPX-JMQ48-75XP herbicide - Use sequential treatments.

With "Puma"

DPX-JMQ48 75XP herbicide can be tank mixed with "Puma" 1EC for control of some annual grass weeds. This tankmix may also include MCP ester, bromoxynil or bromoxynil/MCP, Starane, or Starane + Sword for greater spectrum of broadleaf control - see "Puma" 1EC label for specific use directions and restrictions on tank mixes.

DPX-JMQ48 75XP and 3 to 4 ounces active ingredient per acre of a bromoxynil containing herbicide (such as "Bronate" or "Bison" at 3/4 to 1 pint per acre) may be tank mixed with 0.66 pint per acre of "Puma" for annual grass control in wheat or barley. This tank mix may also include "Starane" for greater spectrum of broadleaf control - see "Puma" label for specific use directions and restrictions. DO NOT use this tank mix on two-row malting barley.

DPX-JMQ48 75XP and a fluroxypyr containing herbicide (such as "Starane" or "Starane +Sword") may be tank mixed with 0.66 pint per acre of "Puma" for annual grass control in wheat or barley. See the "Puma" label for specific use directions, tank mixes, precautions and restrictions of use. This tank mix may also include MCP ester, bromoxynil or bromoxynil/MCP, "Starane", or "Starane + Sword" for greater spectrum of broadleaf control - see "Puma" 1EC label for specific use directions and restrictions on tank mixes. Some reduction in annual grass control may occur when optimum environmental conditions do not occur for several days prior to and after application -- such as low moisture conditions, high and low temperatures, or low humidity.

14/22

With "Tiller"

DuPont™ DPX-JMQ48 75XP can be tank mixed with "Tiller" for green foxtail, foxtail millets and volunteer corn control.

With Other Grass Control Products

DPX-JMQ48 75XP can be tank mixed with grass control products. Antagonism generally does not occur. However, DuPont recommends that you first consult your state experiment station, university, or extension agent, Agricultural dealer, or DuPont representative as to the potential for antagonism before using the mixture. If no information is available, limit the initial use of DPX-JMQ48 75XP and the grass product to a small area.

Do not tank mix DPX-JMQ48 75XP with "Achieve" herbicide".

With Fungicides

DPX-JMQ48 75XP may be tank mixed or used sequentially with fungicides registered for use on cereal grains. Review all fungicide labels for restrictions.

With Insecticides

DPX-JMQ48 75XP may be tank mixed or used sequentially with insecticides registered for use on cereal grains. Review all insecticide labels for restrictions.

However, under certain conditions (drought stress, cold weather, or if the crop is in the 2-4 leaf stage), tank mixes or sequential applications of DPX-JMQ48 75XP with organophosphate insecticides (such as "Lorsban") may produce temporary crop yellowing or, in severe cases, crop injury. The potential for crop injury is greatest when wide fluctuations in day/night temperatures occur just prior to or soon after application. Test these mixtures in a small area before treating large areas.

Do not apply DPX-JMQ48 75XP within 60 days of crop emergence where an organophosphate insecticide has been applied as an in-furrow treatment because crop injury may result.

Do not use DPX-JMQ48 75XP plus "Malathion" because crop injury will result.

With Liquid Nitrogen Solution Fertilizer

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing DPX-JMQ48 75XP in fertilizer solution. DPX-JMQ48 75XP must first be slurried with water and then added to liquid nitrogen solutions (e.g., 28-0-0, 32-0-0).

Ensure that the agitator is running while the DPX-JMQ48 75XP is added. Use of this mixture may result in temporary crop yellowing and stunting.

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of surfactant is necessary. Add surfactant at 1/2 pint - 1 quart per 100 gal of spray solution (0.06 to 0.125% v/v) based on local guidance.

When using high rates of liquid nitrogen fertilizer in the spray solution, adding surfactant increases the risk of crop injury. Consult your agricultural dealer, consultant, fieldsman, or DuPont representative for a specific recommendation before adding an adjuvant to these tank mixtures.

If 2,4-D or MCP is included with an DPX-JMQ48 75XP and fertilizer mixture, ester formulations tend to be more compatible (See manufacturer's label). Additional surfactant may not be needed when using DPX-JMQ48 75XP in tank mix with 2,4-D ester or MCP ester and liquid nitrogen fertilizer solutions. Consult your agricultural dealer, consultant, field advisor, or DuPont representative for a specific recommendation before adding an adjuvant to these tank mixtures.

Liquid nitrogen fertilizer solutions that contain sulfur can increase crop response.

Do not use low rates of liquid fertilizer as a substitute for a surfactant.

Do not use with liquid fertilizer solutions with a pH less than 3.0.

TANK MIXTURES IN FALLOW

DPX-JMQ48 75XP may be used as a fallow treatment, and should be tank mixed with other herbicides that are registered for use in fallow, such as glyphosate (such as Roundup), "Landmaster" II, "Fallow Master", "RT Master", glyphosate plus 2,4-D (ester formulations work best), glyphosate plus dicamba (such as "Banvel"/ "Clarity"), 2,4-D (ester formulations work best), or dicamba (such as "Banvel"/ "Clarity") alone.

DPX-JMQ48 75XP and fluroxypyr containing herbicides (such as "Starane", "Starane +Saber", "Starane +Sword" or "Starane +Salvo") may be used as a fallow treatment, and should be tank mixed with other herbicides that are registered for use in fallow, including glyphosate (such as Roundup), "Landmaster" II, "Fallow Master", "RT Master", glyphosate plus 2,4-D (ester formulations work best), glyphosate plus dicamba (such as "Banvel"/ "Clarity"), 2,4-D (ester formulations work best), or dicamba (such as "Banvel"/ "Clarity") alone.

TANK MIXTURES IN PRE-PLANT BURNDOWN APPLICATIONS

DPX-JMQ48 75XP may be used as a pre-plant burndown treatment alone or tank mixed with other herbicides that are registered for use as a pre-plant burndown product, such as Aim, glyphosate (such as Roundup), "Landmaster" II, "Fallow Master", "RT Master", glyphosate plus dicamba (such as "Banvel"/ "Clarity") or dicamba (such as "Banvel"/ "Clarity") alone.

15/22

TANK MIXTURES IN POST HARVEST APPLICATIONS

DuPont™ DPX-JMQ48 75XP may be used as a post harvest treatment to crop stubble, and should be tank mixed with other herbicides that are registered for use in fallow.

DPX-JMQ48 75XP and fluroxypyr containing herbicides (such as "Starane", "Starane +Saber", "Starane +Sword" or "Starane +Salvo") may be used as a post harvest treatment to crop stubble, and should be tank mixed with other herbicides such as Aim, glyphosate (such as Roundup), "Landmaster" II, "Fallow Master", "RT Master", glyphosate plus dicamba (such as "Banvel" / "Clarity"), or dicamba (such as "Banvel" / "Clarity") alone, that are registered for use in post harvest cereal applications.

FOR BROADLEAF WEED CONTROL IN THE STATE OF ARIZONA

DIRECTIONS FOR USE

DPX-JMQ48 75XP herbicide at 0.4 to 0.7 oz per acre can be used alone or in a tankmix in the state of Arizona for control of broadleaf weeds in wheat, barley, oat and triticale. It is a violation of federal law to use this product in a manner inconsistent with its labeling.

SURFACTANTS

Include a spray adjuvant with applications of DPX-JMQ48 75XP. An ammonium nitrogen fertilizer may also be used. Do not use low rates of liquid nitrogen fertilizer solution as a substitute for a surfactant. Antifoaming agents may be used if needed. Consult the DPX-JMQ48 75XP product label for specific adjuvant recommendations.

WEEDS CONTROLLED

- Annual sowthistle
- Black mustard
- Bushy wallflower/ Treacle mustard
- Coast fiddleneck
- Common chickweed*
- Common groundsel
- Common lambsquarters
- Curly dock
- Kochia *†
- London rocket
- Mallow (little)
- Miners-lettuce
- Prostrate knotweed
- Redmaids
- Redroot pigweed†
- Russian thistle*†
- Shepherd's-purse
- Stinking mayweed/Dogfennel
- Swinecress
- Tumble/Jim Hill mustard
- Wild mustard†

WEEDS PARTIALLY CONTROLLED**

- Common cocklebur†
- Common sunflower†
- Cutleaf eveningprimrose
- Deadnettle (purple, red)
- Henbit
- Mallow (common)
- Prickly lettuce*†
- Tansymustard*
- Wild radish*

* See "SPECIFIC WEED INSTRUCTIONS" for more information.

**Partial control: A visual reduction of weed population as well as a significant loss of vigor for individual weed plants. For better results, use 0.5 to 0.7 ounce DPX-JMQ48 75XP per acre and include a tank mix partner (refer to the "TANK MIXTURES" section of the product label).

† Naturally occurring resistant biotypes are known to occur.

SPECIFIC WEED INSTRUCTIONS

Common chickweed: For best results, apply a minimum of 0.5 ounce DPX-JMQ48 75XP per acre plus surfactant when all or the majority of weeds have germinated and are past the cotyledon stage. Weeds should be less than 3 inches tall or across at the time of DPX-JMQ48 75XP application.

Kochia: Naturally occurring biotypes resistant to DPX-JMQ48 75XP are known to occur. For best results, use DPX-JMQ48 75XP in a tank mix with "Starane", "Starane + Salvo", "Starane + Sword", dicamba (such as "Banvel" / "Clarity") and 2,4-D or MCP (ester or amine), or bromoxynil containing products (such as "Buctril", "Bison", "Bronate" or "Bronate Advanced"). DPX-JMQ48 75XP should be applied in the spring when kochia are less than 2" tall and are actively growing (refer to the "TANK MIXTURES" section of the label for additional details on rates and restrictions).

Russian thistle, Prickly lettuce: Naturally occurring biotypes resistant to DPX-JMQ48 75XP of these weeds are known to occur. For best results, use DPX-JMQ48 75XP in a tank mix with dicamba (such as "Banvel" / "Clarity") and 2,4-D or MCP (ester or amine), or bromoxynil containing products (such as "Buctril", "Bison", "Bronate" or "Bronate Advanced").

16/22

DuPont™ DPX-JMQ48 75XP should be applied in the spring when Russian thistle and prickly lettuce are less than 2" tall or 2" across and are actively growing (refer to the "TANK MIXTURES" section of the label for additional details on rates and restriction).

Wild radish: For best results, apply 0.5 to 0.7 ounce DPX-JMQ48 75XP per acre plus surfactant either in the fall or spring to wild radish rosettes less than 6 inches in diameter. Applications made later than 30 days after weed emergence will result in partial control. Fall applications should be made prior to hardening-off of plants.

GRAZING

Allow at least 7 days between application and grazing of treated forage. In addition, allow at least 7 days between application and feeding of forage from treated areas to livestock. Allow at least 30 days between application and feeding of hay from treated areas to livestock. Harvested straw may be used for bedding and/or feed. Allow at least 45 days between application and harvesting of grain.

CROP ROTATION

Labeled crops may be planted at specified time intervals following application of labeled rates of DPX-JMQ48 75XP. Use the time intervals listed below to determine the required time interval before planting.

Time Interval Before Planting*
(days after treatment with DPX-JMQ48 75XP)

Crop	Days
Barley, Rice, Triticale, and Wheat (including durum)	0
Oat and Soybeans	1**
Cotton, Field Corn, and Grain/forage Sorghum	14**
Sugarbeets, Winter Rape, and Canola	60
Any other crop	45

* Refer to individual product labels to determine rotational crop restrictions when tank mixtures are used.

**Where DPX-JMQ48 75XP is used on light textured soils, such as sands and loamy sands, extend time to planting by 7 additional days. Where DPX-JMQ48 75XP is used on high-pH soils (>7.9), extend time to planting by 7 additional days.

APPLICATION INFORMATION

PRODUCT MEASUREMENT

DPX-JMQ48 75XP can be measured using the DPX-JMQ48 75XP volumetric measuring cylinder provided by DuPont. The degree of accuracy of this cylinder varies by +/- 7.5%. For more precise measurement, use scales calibrated in ounces.

MIXING INSTRUCTIONS

Do not use with spray additives that alter the pH of the spray solution below pH 5.0 or above pH 9.0, as rapid product degradation can occur. Spray solutions of pH 6.0 - 8.0 allow for optimum stability of DPX-JMQ48 75XP.

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of DPX-JMQ48 75XP.
3. Continue agitation until the DPX-JMQ48 75XP is fully dispersed, at least 5 minutes.
4. Once the DPX-JMQ48 75XP is fully dispersed, maintain agitation and continue filling tank with water. DPX-JMQ48 75XP should be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the required volume of spray adjuvant. Always add spray adjuvant last. Antifoaming agents may be used. Do not use with spray additives that alter the pH of the spray solution below pH 6.0 as rapid product degradation can occur. Spray solutions of pH 7.0 and higher allow for optimum stability of DPX-JMQ48 75XP.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply DPX-JMQ48 75XP spray mixture within 24 hours of mixing to avoid product degradation.
8. If DPX-JMQ48 75XP and a tank mix partner are to be applied in multiple loads, pre-slurry the DPX-JMQ48 75XP in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the DPX-JMQ48 75XP.

GROUND APPLICATION

For optimum spray distribution and thorough coverage, use flat-fan or low-volume flood nozzles.

- For best performance, select nozzles and pressure that deliver MEDIUM spray droplets.
- Nozzles that deliver COARSE spray droplets may be used to reduce drift, provided spray volume is increased to maintain coverage on small weeds. For optimal product performance and minimal spray drift, adjust the spray boom to the lowest possible spray height recommended in manufacturers' specifications.
- Overlaps or starting, stopping, slowing, and turning while spraying may result in crop injury.

- For flat-fan nozzles, use a spray volume of at least 5 gal per acre (GPA).
- For flood nozzles on 30" spacings, use at least 10 GPA, flood nozzles no larger than TK10 (or the equivalent), and a pressure of at least 30 psi. For 40" nozzle spacings, use at least 13 GPA; for 60" spacings use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.
- "Raindrop RA" nozzles are not recommended for DuPont™ DPX-JMQ48 75XP herbicide applications, as weed control performance may be reduced.
- Use screens that are 50-mesh or larger.

AERIAL APPLICATION

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage.

- Use 2 to 5 GPA
- Use at least 3 GPA in Idaho, Oregon, or Utah

Do not apply DPX-JMQ48 75XP by air in the state of New York.

When applying DPX-JMQ48 75XP by air in areas adjacent to sensitive crops, use solid stream nozzles oriented straight back. Adjust the swath to avoid spray drift damage to sensitive crops downwind and/or use ground equipment to treat the border edge of fields. See the "SPRAY DRIFT MANAGEMENT" section of this label.

SPRAY EQUIPMENT

For specific application equipment, refer to the manufacturer's instructions for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc.

Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when crop canopy is dense. Avoid swath overlapping, and shut off spray booms while starting, turning, slowing, or stopping, to avoid injury to the crop. Do not make applications using equipment and/or spray volumes or during weather conditions that might cause spray to drift onto nontarget sites. For additional information on spray drift refer to the "SPRAY DRIFT MANAGEMENT" section of this label.

Continuous agitation is required for DPX-JMQ48 75XP to keep tank-mix partners in solution or suspension. Refer to tank-mix partner labels for additional information.

SPRAYER CLEANUP

The spray equipment must be cleaned before DPX-JMQ48 75XP is sprayed. Follow the cleanup procedures specified on the labels of the previously applied products. If no directions are provided, follow the six steps outlined in the "AFTER SPRAYING DPX-JMQ48 75XP" section of this label.

AT THE END OF THE DAY

It is recommended that during periods when multiple loads of DPX-JMQ48 75XP herbicide are applied, at the end of each day of spraying the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits which can accumulate in the application equipment.

AFTER SPRAYING DPX-JMQ48 75XP AND BEFORE SPRAYING CROPS OTHER THAN WHEAT, BARLEY, OAT, TRITICALE, FIELD CORN AND SOYBEANS

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of DPX-JMQ48 75XP as follows:

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and 1 gal of household ammonia* (contains 3% active ingredient) for every 100 gal of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 min. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat step 2.
5. Rinse the tank, boom, and hoses with clean water.
6. If only ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) specified on this label. Do not exceed the maximum labeled use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

* Equivalent amounts of an alternate-strength ammonia solution or a DuPont-approved cleaner can be used in the cleanup procedure. Carefully read and follow the individual cleaner instructions. Consult your Ag dealer, applicator, or DuPont representative for a listing of approved cleaners.

Notes:

1. **CAUTION:** Do not use chlorine bleach with ammonia because dangerous gases will form. Do not clean equipment in an enclosed area.
2. Steam-cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.
3. When DuPont™ DPX-JMQ48 75XP herbicide is tank mixed with other pesticides, all cleanout procedures for each product should be examined and the most rigorous procedure should be followed.
4. In addition to this cleanout procedure, all precleanout guidelines on subsequently applied products should be followed as per the individual product labels.
5. Where routine spraying practices include shared equipment frequently being switched between applications of DPX-JMQ48 75XP and applications of other pesticides to DPX-JMQ48 75XP-sensitive crops during the same spray season, it is recommended that a sprayer be dedicated to DPX-JMQ48 75XP to further reduce the chance of crop injury.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions. **AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.**

IMPORTANCE OF DROPLET SIZE

The most effective drift management strategy is to apply the largest droplets which are consistent with pest control objectives. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!**

A droplet size classification system describes the range of droplet sizes produced by spray nozzles. The American Society of Agricultural and Biological Engineers (ASABE) provide a Standard that describes droplet size spectrum categories defined by a number of reference nozzles (fine, coarse, etc.). Droplet spectra resulting from the use of a specific nozzle may also be described in terms of volume mean diameter (VMD). Coarser droplet size spectra have larger VMD's and lower drift potential.

CONTROLLING DROPLET SIZE - GROUND APPLICATION

- **Nozzle Type** - Select a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. The use of low-drift nozzles will reduce drift potential.
- **Pressure** - The lowest spray pressures recommended for the nozzle produce the largest droplets. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, using a higher-capacity nozzle instead of increasing pressure results in the coarsest droplet spectrum.
- **Flow Rate/Orifice Size** - Using the highest flow rate nozzles (largest orifice) that are consistent with pest control objectives reduces the potential for spray drift. Nozzles with higher rated flows produce coarser droplet spectra.

CONTROLLING DROPLET SIZE – AIRCRAFT

- **Nozzle Type** - Solid stream, or other low drift nozzles produce the coarsest droplet spectra.
- **Number of Nozzles** - Using the minimum number of nozzles with the highest flow rate that provide uniform coverage will produce a coarser droplet spectrum.
- **Nozzle Orientation** - Orienting nozzles in a manner that minimizes the effects of air shear will produce the coarsest droplet spectra. For some nozzles such as solid stream, pointing the nozzles straight back parallel to the airstream will produce a coarser droplet spectrum than other orientations.
- **Pressure** - Selecting the pressure that produces the coarsest droplet spectrum for a particular nozzle and airspeed reduces spray drift potential. For some nozzle types such as solid-streams, lower pressures can produce finer droplet spectra and increase drift.

BOOM LENGTH (AIRCRAFT), AND APPLICATION HEIGHT

- **Boom Length (aircraft)** - Using shorter booms decreases drift potential. Boom lengths are expressed as a percentage of an aircraft's wingspan or a helicopter's rotor blade diameter. Shorter boom length and proper positioning can minimize drift caused by wingtip or rotor vortices.
- **Application Height (aircraft)** - Applications made at the lowest height that are consistent with pest control objectives and the safe operation of the aircraft will reduce the potential for spray drift.
- **Application Height (ground)** - Applications made at the lowest height consistent with pest control objectives, and that allow the applicator to keep the boom level with the application site and minimize bounce, will reduce the exposure of spray droplets to evaporation and wind, and reduce spray drift potential.

WIND

Drift potential is lowest when applications are made in light to gentle sustained winds (2-10 mph), which are blowing in a constant direction. Many factors, including droplet size and equipment type also determine drift potential at any given wind speed. **AVOID GUSTY OR WINDLESS CONDITIONS.**

Local terrain can also influence wind patterns. Every applicator is expected to be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

Setting up equipment to produce larger droplets to compensate for droplet evaporation can reduce spray drift potential. Droplet evaporation is most severe when conditions are both hot and dry.

SURFACE TEMPERATURE INVERSIONS

Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which may cause small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Mist or fog may indicate the presence of an inversion in humid areas.

Inversions may also be identified by producing smoke and observing its behavior. Smoke that remains close to the ground, or moves laterally in a concentrated cloud under low wind conditions indicates a surface inversion. Smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, that it is configured properly, and that drift potential has been minimized.

Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Read the specific crop use and application equipment instructions to determine if an air assisted field crop sprayer can be used.

SENSITIVE AREAS

Making applications when there is a sustained wind moving away from adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is an effective way to minimize the effect of spray drift.

DRIFT CONTROL ADDITIVES

Using product compatible drift control additives can reduce drift potential. When a drift control additive is used, read and carefully observe cautionary statements and all other information on the additive's label. If using an additive that increases viscosity, ensure that the nozzles and other application equipment will function properly with a viscous spray solution. Preferred drift control additives have been certified by the Chemical Producers and Distributors Association (CPDA).

20/22

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store product in original container only. Store in a cool, dry place.

Pesticide Disposal: Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities

Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums With Liners: Nonrefillable container. Do not reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

Refillable Fiber Drums With Liners: Refillable container (fiber drum only). *Refilling Fiber Drum:* Refill this fiber drum with DuPont™ DPX-JMQ48 75XP herbicide containing thifensulfuron methyl and tribenuron methyl only. Do not reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. *Disposing of Fiber Drum and/or Liner:* Do not reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

All Other Refillable Containers: Refillable container. *Refilling Container:* Refill this container with DuPont™ DPX-JMQ48 75XP herbicide containing thifensulfuron methyl and tribenuron methyl only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use the container, contact DuPont at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found; do not reuse or transport container, contact DuPont at the number below for instructions. *Disposing of Container:* Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Outer Foil Pouches of Water Soluble Packets (WSP): Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or, dispose of the empty outer foil pouch in the trash as long as WSP is unbroken. If the outer pouch contacts the formulated product in any way, the pouch must be triple rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer pouch as described previously.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact DuPont at 1-800-441-3637, day or night.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

The DuPont Oval Logo and DuPont™ are trademarks or registered trademarks of E. I. duPont de Nemours & Company
"Maverick", "Fallow Master", "RT Master" and "Roundup" are trademarks or registered trademarks of Monsanto Technology LLC

"Assert" and "Weedar" are trademarks or registered trademarks of NuFarm Americas, Inc.

"Banvel", "Clarity" and "Clearfield" are trademarks or registered trademarks of BASF Corporation.

"Puma", "Buctril", "Bronate", "Bronate Advanced", "Tiller", "Huskie", "Wolverine" and "Hoelon" are trademarks or registered trademarks of Bayer CropScience

"Rhino" is a registered trademark of The McGreagor Company

"Everest" is a registered trademark of Arvesta LifeScience North America LLC

"Aim" is a registered trademark of FMC Corporation

"Achieve", "Axial" and "Discover" are registered trademarks of Syngenta Crop Protection

"CleanWave", "Stinger", "Curtail", "Curtail M", "Lorsban", "Starane", and "Widematch" are trademarks or registered trademarks of Dow AgroSciences LLC

"Landmaster" is a registered trademark of Albaugh, Inc

"Colt", "Salvo" and "Sword" are trademarks or registered trademarks of Loveland Products, Inc.

"Bison" is a registered trademark of Winfield Solutions LLC

"Barrage" is a registered trademark of Helena Holding Company

D - 1817MSTR 073114

**LIMITATION OF
WARRANTY AND LIABILITY**

NOTICE: Read this Limitation of Warranty and Liability Before Buying or Using This Product. DuPont Will Not Be Responsible for Losses or Damages Resulting from the Use of This Product in Any Manner Not Specifically Directed by DuPont. User Assumes All Risks Associated With Such Non-Directed Use.

If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of DuPont. These risks can cause: ineffectiveness of the product, crop injury, or injury to non-target crops or plants. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

DuPont warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, DUPONT MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL DUPONT OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF DUPONT OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF DUPONT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

To the extent consistent with applicable law that allows such requirement, DuPont or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify DuPont or a DuPont Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise, or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

For product information call: 1-888-6-DUPONT [1-888-638-7668]

Internet address: <http://cropprotection.dupont.com/>

© 2014 E. I. du Pont de Nemours and Company, 1007 Market Street, Wilmington, DE 19898

All rights reserved.